

- 10 -

Claims:

1. A fish feed comprising a live feed component which has been fed with an acylglycerol composition which
5 comprises mono and/or diacylglycerols of at least one fatty acid selected from eicosapentaeneoic acid and docosaheptaeneoic acid.
2. A fish feed as claimed in claim 1 wherein the
10 content of said mono and/or diacylglycerols is at least 45% wt of the total acylglycerol content.
3. A fish feed as claimed in either one of claims 1 or 2 wherein said fish are flatfish larvae.
- 15 4. A process for the preparation of fish feed, said process comprising contacting an acylglycerol composition and a live feed component characterised in that said acylglycerol composition comprises mono and/or
20 diacylglycerols of at least one fatty acid selected from eicosapentaeneoic acid and docosaheptaeneoic acid.
5. A process as claimed in claim 4 wherein the content of said mono and/or diacylglycerols is at least 45% wt
25 of the total acylglycerol content.
6. A process as claimed in either one of claims 4 or 5 wherein said acylglycerol composition comprises at least mono and diacylglycerols containing EPA and/or DHA
30 residues.
7. A process as claimed in any one of claims 4 to 6 wherein said acylglycerol composition contains
acylglycerols having arachidonic acid (AA) residues.
35
8. A process as claimed in any one of claims 4 to 7 wherein said acylglycerol is fed to said live feed

- 11 -

component at a concentration of about 0.05 to 0.10 g/l.

9. The use in the preparation of a fish feed, e.g. a formulated fish feed, of mono and/or diacylglycerols containing EPA and/or DHA residues.

10. A method of raising fish comprising feeding fish or fish larvae with an acylglycerol-containing feed, characterized in that said acylglycerol comprises mono and/or diacylglycerols containing EPA and/or DHA residues.

11. A method of raising fish comprising feeding filter-feeding fish with an acylglycerol containing feed, characterized in that said feed comprises an oil-in-water emulsion containing mono and/or diacylglycerols of at least one fatty acid selected from EPA and DHA.

12. An aquatic animal organism having a maximum dimension of 2mm and having in its digestive tracts mono and/or diacylglycerols containing EPA and/or DHA residues.

13. Milk powder containing mono and/or diacylglycerols containing EPA and/or DHA residues.

14. Puréed cooked fruit, meat or vegetable containing mono and/or diacylglycerols containing EPA and/or DHA residues.

15. A diacylglycerol wherein both fatty acid groups are EPA, DHA and/or AA residues.